



ЭКОЛОГИЯ/ECOLOGY

DOI: <https://doi.org/10.60797/JAE.2026.74.13> EDN: YYMRQD**PLANORBIDAE SNAIL EMBRYOS AS A TEST OBJECT FOR ECOLOGICAL MONITORING OF LITHIUM POLLUTION IN AQUATIC BODIES**

Research article

Kuzyanov D.A.^{1,*}, Moiseeva E.M.², Erdniev L.P.³¹ORCID : 0000-0002-5070-4431;^{1,2,3} Saratov Medical Research Center for Hygiene, a branch of the Federal Scientific Center for Medical and Preventive Technologies for Public Health Risk Management, Saratov, Russian Federation

* Corresponding author (dimakuzyanov2000[at]gmail.com)

Suggested: 26.08.2026; Accepted: 30.09.2026; Published: 19.10.2026

Abstract

The degradation of water resource quality and ecosystem decline driven by increasing lithium contamination represents a major global challenge, necessitating the search for potential biological indicators to assess the ecological status of water bodies. The objective of this study was to evaluate the suitability of gastropod embryos from the *Planorbidae* family as a test object under conditions of aquatic lithium pollution. Experimental modeling of the effects of a lithium compound on *Planorbidae* embryos established a clear relationship between element concentration and changes in embryo size during ontogenesis; furthermore, the effective concentrations were compared against the maximum permissible concentrations (MPC) established for fishery water bodies. The EC₂₀ value for lithium was 0.07 mg/L (0.9 of the MPC), while the EC₅₀ was 0.28 mg/L (exceeding the MPC by 3.5 times). These results demonstrate that *Planorbidae* embryos hold significant promise for assessing the ecological status of aquatic environments affected by lithium pollution.

Keywords: aquatic ecological monitoring, embryotoxicity, lithium contamination, gastropods, Planorbidae embryos.**ЭМБРИОНЫ УЛИТОК PLANORBIDAE КАК ТЕСТ-ОБЪЕКТ В ЭКОЛОГИЧЕСКОМ КОНТРОЛЕ ЛИТИЕВОГО ЗАГРЯЗНЕНИЯ ВОДНЫХ ОБЪЕКТОВ**

Научная статья

Кузянов Д.А.^{1,*}, Моисеева Е.М.², Эрднеев Л.П.³¹ORCID : 0000-0002-5070-4431;^{1,2,3} Саратовский медицинский научный центр гигиены, филиал Федерального научного центра медико-профилактических технологий управления рисками здоровью населения, Саратов, Российская Федерация

* Корреспондирующий автор (dimakuzyanov2000[at]gmail.com)

Предложена: 26.08.2026; Принята: 30.09.2026; Опубликована: 19.10.2026

Аннотация

Проблема ухудшения качества водных ресурсов и деградации экосистем вследствие нарастающего загрязнения соединениями лития является одной из наиболее значимых в мировом масштабе, что определяет необходимость поиска потенциальных биологических объектов для оценки экологического состояния водоемов. Цель настоящего исследования заключалась в определении возможности использования эмбрионов брюхоногих моллюсков семейства *Planorbidae* в качестве тест-объекта в условиях загрязнения водных объектов соединениями лития. Экспериментальное моделирование воздействия соединения лития на эмбрионы моллюсков *Planorbidae* позволило установить зависимость между концентрацией элемента и изменением размера эмбрионов в процессе онтогенеза, а также сопоставить его эффективные концентрации нормативам содержания в водных объектах рыбохозяйственного значения. Величина ЕС₂₀ лития составила 0,07 мг/л (0,9 от ПДК), а ЕС₅₀ — 0,28 мг/л (превышение ПДК в 3,5 раза). Результаты демонстрируют перспективность использования эмбрионов моллюсков *Planorbidae* для оценки экологического состояния водных объектов в условиях их загрязнения литием.

Ключевые слова: экологический контроль водных объектов, эмбриотоксичность, загрязнение литием, брюхоногие моллюски, эмбрионы Planorbidae.**Introduction**

Along with persistent organic pollutants, nutrients, and xenobiotics, compounds of alkali and alkaline earth metals—specifically lithium—are among the most significant water contaminants. Lithium is widely utilized across the glass, chemical, metallurgical, and medical industries. Chemical power sources account for more than half of global lithium consumption, and by 2050, demand for this element is projected to rise significantly, which will inevitably intensify mining activities and consequently drive environmental pollution. This stems from the fact that the primary entry pathways of lithium into water bodies are associated with enterprises involved in lithium ore extraction, processing, and lithium-ion battery manufacturing. Furthermore, due to low recycling rates of lithium-containing products, landfill leachate from municipal solid waste sites also acts as a notable source of contamination. Concurrently, natural geological processes further contribute to the accumulation of this element in aquatic environments [1].

The maximum permissible concentration (MPC) of lithium in fishery water bodies, established to prevent negative impacts on aquatic biological resources and their habitats, is 0.08 mg/L. Meanwhile, lithium concentrations in water bodies vary across a wide range. The lowest lithium levels, ranging from 0.0009 to 0.0099 mg/L, are recorded in major river systems (e.g., the Volga and Tom rivers), with an increasing concentration trend driven by urban runoff [3]. Concurrently, elevated lithium levels are consistently observed in aquatic bodies within developed geological and industrial regions, as well as in tectonic fault zones. For instance, in the Lugan River, located within an active coal-mining area, the lithium concentration reaches 0.14 mg/L. Discharges from mine drainage and processing plants, along with runoff from waste heaps (where lithium content reaches up to 5.59 mg/L), maintain lithium levels in the Lozovaya River at 0.09 mg/L, and in the Kamyshevskaya River at up to 0.08 mg/L [4]. The water of the Lower Don basin and the Mius River is characterized by lithium concentrations up to 0.145 mg/L, which is attributed to industrial wastewater inputs [5]. Lithium contamination in the Ardon and Sadon river basins (0.066–0.068 mg/L) is associated with mining activities at the lead-zinc ore deposits of the Sadon mining district [6].

The high mobility of the lithium ion in water and its resistance to complexation render the element highly bioavailable to aquatic organisms. The toxic effects of lithium are known to manifest as alterations in the physiological and behavioral responses of living organisms [7]. Consequently, the contamination of water bodies with lithium compounds can adversely affect the stability of freshwater ecosystems. However, ecotoxicological data regarding its impact on aquatic life remain limited [7]. The embryonic and larval stages of aquatic organisms, particularly mollusks, are among the most sensitive to lithium exposure [8]. Toxic effects during the embryonic stages of mollusks can reduce the population of adult individuals involved in nutrient cycling and water self-purification, subsequently disrupting the ecological balance. This underscores the relevance and urgency of utilizing these biological test systems to assess the ecological status of lithium-polluted water bodies.

The objective of this study was to evaluate the suitability of gastropod embryos from the *Planorbidae* family as a test object under conditions of aquatic lithium pollution.

Research methods and principles

The objects of this study were gastropods of the *Planorbidae* family, which were collected from a natural population in an unpolluted water body located within the Saratov Oblast and subsequently acclimated to laboratory conditions.

The mollusks were maintained in a straight-walled transparent container filled with tap water that had been aged for 5 days, at a temperature of 20°C and a pH range of 6.6–7.2.

To evaluate the sensitivity of the *Planorbidae* mollusks, synchronized egg capsules at the zygote stage (at least 3 capsules per replicate) were placed into 2-mL tubes, to which 1 mL of an aqueous lithium sulfate solution (Li_2SO_4), analytical grade, JSC Vekton, St. Petersburg, was added. The experimental concentrations, calculated as lithium ion equivalent, were 0.05, 0.1, 1, and 10 mg/L (treatment groups). Each synchronized egg mass contained at least 6 embryos. Aged tap water was used as the control. Both control and treatment groups were exposed for 7 days from the initial contact of the egg capsules with the solution, maintaining a constant temperature of 20°C.

The test response used to evaluate the toxic impact of the lithium compound was the change in the size of the mollusk embryos within the egg masses in the treatment solutions compared to the control after 7 days of exposure (embryotoxicity); the relative change in embryo size was determined according to Equation 1:

$$\text{Embryotoxicity} = \frac{B-A}{B} \times 100\%$$

where:

A is the mean size of the embryos within the egg mass in the treatment group;

B is the mean size of the embryos within the egg mass in the control group.

Embryo sizes were measured every 24 hours throughout the exposure period using light microscopy (up to 12.5× magnification) coupled with a digital camera (ToupCam UA1600CA) for image capture overlaid with a grid field (100 µm cell size). The size of the embryos was determined by calculating their surface area (mm²) using the ImageJ software (version 1.42, Fiji bundle).

Statistical data processing was performed using the StatTech software, version 4.5.0 (Stattech LLC, Russia). In the statistical analysis of the results, mean values and standard deviations were evaluated. The calculation of effective concentrations (EC_{20} and EC_{50}) along with their 95% confidence intervals was carried out using GraphPad Prism software (GraphPad Software, USA), employing a regression model based on a four-parameter logistic approximation.

Main results and discussion

Water biotesting using gastropod mollusks (Gastropoda) can serve as a promising tool for identifying the negative impacts of contaminants—specifically lithium—on aquatic ecosystems due to the ecological relevance of these test objects. Within the taxonomic class Gastropoda, the *Planorbidae* family is the largest and most promising for water toxicity studies owing to a relatively short embryo incubation period (one to two weeks), flat egg masses suitable for microphotography, and a high egg production capacity [9]. Mollusks of the *Planorbidae* family are typical representatives of freshwater fauna whose embryonic and larval developmental stages occur entirely within the aquatic environment, rendering them highly susceptible to chemical compounds present in this medium [9].

In the present study, to determine the sensitivity of *Planorbidae* embryos to lithium pollution, the impact of the element at various concentrations on egg masses was evaluated in a model experiment. A dose-dependent effect of lithium exposure at concentrations of 0.05–10 mg/L was established, manifesting as the inhibition of embryonic development. The dynamics of changes in *Planorbidae* embryo sizes under lithium exposure are presented in Figure 1.




Figure 1 - Dynamics of changes in *Planorbidae* embryo sizes under lithium exposure throughout the experiment
DOI: <https://doi.org/10.60797/JAE.2026.74.13.1>

According to the obtained results, the size of the egg capsules in the control group increased 35-fold by day 7 of the experiment compared to day 1. Under lithium exposure at the lowest investigated concentration (0.05 mg/L), the size of the egg capsules during the first 4 days was statistically significantly larger than the corresponding values in the control group. This is likely attributed to the effect of hormesis—the stimulating effect of low concentrations of chemical substances on biological systems [10]. However, starting from the fifth day of exposure, the intensity of embryonic growth began to decline, resulting in a 1.4-fold reduction in capsule size relative to the control group.

Growth inhibition in the mollusks exposed to lithium at concentrations of 0.1, 1.0, and 10.0 mg/L was recorded as early as day 2 of exposure. After 7 days of exposure, the size of the egg capsules treated with lithium at a concentration of 0.1 mg/L was 1.77-fold lower than that of the control group, and at a concentration of 1.0 mg/L, it was 4.88-fold lower. The most pronounced effect of lithium was observed at a concentration of 10 mg/L; by the end of the experiment, the size of the control capsules exceeded that of the exposed ones 25-fold.

The obtained results demonstrate the sensitivity of mollusk embryonic stages to lithium exposure and are consistent with data from earlier international studies. In particular, it has been shown that lithium exposure at concentrations starting from 0.015 mg/L can influence the growth of *Marisa cornuarietis* embryos [8]. Conversely, other studies on *Mytilus galloprovincialis* have reported a decrease in acetylcholinesterase activity and impairment of the antioxidant defense system in the presence of 0.1 mg/L of lithium, as well as an increased rate of burrowing into the bottom substrate in the gastropod *Tritia reticulata* at 0.5 mg/L [11].

To evaluate the trend of the embryotoxic effect of lithium at the investigated concentrations on the mollusk egg masses, the ratios of the mean embryo sizes in the treatment solutions to those in the control group were assessed on days 1 and 7 of exposure, and a cumulative dose-response curve was constructed (Figure 2).




Figure 2 - Dose-response curve for the effects of lithium ions on *Planorbidae* embryos

DOI: <https://doi.org/10.60797/JAE.2026.74.13.2>

The application of a probit regression model enabled the interpolation of effective lithium concentrations causing 20% (EC_{20}) and 50% (EC_{50}) reductions in embryo size, which were determined to be 0.07 mg/L and 0.28 mg/L, respectively. Importantly, the calculated EC_{20} value was virtually at the MPC level established for fishery water bodies, representing 0.9 of the actual standard, whereas the EC_{50} value exceeded the MPC by a factor of 3.5.

The values of the model's coefficient of determination ($R^2 = 0.842$) and the chi-squared statistic ($X^2 = 12.45$), $p = 0.002$ for the probit model indicate a high degree of goodness-of-fit to the experimental data and confirm the adequacy of the obtained results.

In conclusion, the obtained results demonstrate the feasibility of utilizing *Planorbidae* embryo cultures as a test object to evaluate the impact of aquatic lithium pollution on the embryonic developmental stages of aquatic organisms. The findings of the present study can be further applied to the development and standardization of a biotesting method using mollusks of this family. Additionally, assessing the potential of *Planorbidae* mollusks as bioindicators of aquatic lithium contamination is of significant interest. The high sensitivity of these organisms to the toxicant and its inhibitory effect on growth processes indicate that under conditions of lithium pollution in a water body, the population may undergo degradation or complete extinction. Conducting experiments on real-world samples under conditions of lithium contamination represents a promising avenue for further research.

Conclusion

Based on the results of the conducted study, the feasibility of using gastropod embryos of the *Planorbidae* family as a test object under conditions of aquatic lithium pollution was determined. A dose-dependent pattern of lithium toxic action was established, and the effective concentrations of the element were compared against the MPC established for fishery water bodies: the EC_{20} value was 0.07 mg/L (0.9 of the MPC) and the EC_{50} value was 0.28 mg/L (exceeding the MPC by a factor of 3.5). Based on the evaluation of the ecological relevance of the test object to conditions characteristic of the aquatic environment, the determination of its sensitivity to lithium, the analysis of the test response, and the alignment of the experimental modeling results with the established standards for the element content in fishery water bodies, the use of these mollusk embryos holds significant promise for assessing the ecological status of aquatic environments affected by lithium pollution.

Конфликт интересов

Не указан.

Рецензия

Все статьи проходят рецензирование. Но рецензент или автор статьи предпочли не публиковать рецензию к этой статье в открытом доступе. Рецензия может быть предоставлена компетентным органам по запросу.

Conflict of Interest

None declared.

Review

All articles are peer-reviewed. But the reviewer or the author of the article chose not to publish a review of this article in the public domain. The review can be provided to the competent authorities upon request.



Список литературы / References

1. Ruseva A.S. Assessment of Lithium Ecotoxicity in Soils with Contrasting Properties / A.S. Ruseva, N.A. Evstegneeva, S.I. Kolesnikov et al. // Eurasian Soil Science. — 2025. — № 58 (10). — P. 136. — DOI: 10.1134/S1064229325601106
2. Об утверждении нормативов качества воды водных объектов рыбохозяйственного значения, в том числе нормативов предельно допустимых концентраций загрязняющих веществ в водах водных объектов рыбохозяйственного значения : Приказ Федерального агентства по рыболовству от 26 мая 2025 г. № 296 // Официальный интернет-портал правовой информации. — 2025. — URL: <http://publication.pravo.gov.ru/document/0001202506020069>. (дата обращения: 24.08.26)
3. Сазонов А. А. Синхронные исследования качества воды реки Волги / А. А. Сазонов, А. А. Лисина, О. Н. Ерина [и др.] // Водное хозяйство России: проблемы, технологии, управление. — 2024. — № 5. — С. 98–114. — DOI: 10.35567/19994508-2024-5-98-114
4. Удалов И. В. Гидрохимическая характеристика поверхностных и грунтовых вод Лисичанского и Алмазно-Марьевского геолого-промышленных районов Северо-Восточного Донбасса / И. В. Удалов // Journal of Geology, Geography and Geoecology. — 2014. — Т. 22. — № 3 (2). — С. 2–11.
5. Хорошевская В. О. Сравнительная характеристика микроэлементного состава речных вод, дренирующих Донецкий и Кузбасский угольные бассейны / В. О. Хорошевская // Международный научно-исследовательский журнал. — 2022. — № 5-1 (119). — С. 124–134. — DOI: 10.23670/IRJ.2022.119.5.062
6. Реутова Н. В. Природное и антропогенное загрязнение микроэлементами верховий бассейна р. Ардон / Н. В. Реутова, Т. В. Реутова, Ф. Р. Дреева // Геология и геофизика Юга России. — 2025. — Т. 15. — № 2. — С. 219–230. — DOI: 10.46698/VNC.2025.46.62.001
7. Batley G. E. Metal contaminants of emerging concern in aquatic systems / G. E. Batley, P. G. C. Campbell // Environmental Chemistry. — 2022. — Vol. 19. — № 1. — P. 23–40. — DOI: 10.1071/EN22030
8. Sawasdee B. Metal sensitivity of the embryonic development of the ramshorn snail *Marisa cornuarietis* (Prosobranchia) / B. Sawasdee, H. R. Köhler // Ecotoxicology. — 2010. — Vol. 19. — № 8. — P. 1487–1495. — DOI: 10.1007/s10646-010-0534-8
9. Osborne R. K. Development of an embryo toxicity test to assess the comparative toxicity of metal exposure on different life stages of freshwater gastropods / R. K. Osborne, M. M. Ijzerman, C. Venier [et al.] // Environmental Toxicology and Chemistry. — 2023. — Vol. 42. — № 8. — P. 1791–1805. — DOI: 10.1002/etc.5686
10. Calabrese E. J. Lithium and hormesis: Enhancement of adaptive responses and biological performance via hormetic mechanisms / E. J. Calabrese, P. Pressman, A. W. Hayes [et al.] // Journal of Trace Elements in Medicine and Biology. — 2023. — Vol. 78. — P. 127156. — DOI: 10.1016/j.jtemb.2023.127156
11. Noguera L. Integration of responses to lithium in mussels at different levels of biological complexity / L. Noguera, D. Benito, T. Briaudeau [et al.] // Chemosphere. — 2025. — Vol. 371. — P. 144021. — DOI: 10.1016/j.chemosphere.2024.144021.
12. Campos S. Behavioural and biochemical responses of the sea snail *Tritia reticulata* to lithium concentration gradient / S. Campos, C. Leite, J. Pinto [et al.] // Aquatic Toxicology. — 2023. — Vol. 261. — P. 106629. — DOI: 10.1016/j.aquatox.2023.106629.

Список литературы на английском языке / References in English

1. Ruseva A.S. Assessment of Lithium Ecotoxicity in Soils with Contrasting Properties / A.S. Ruseva, N.A. Evstegneeva, S.I. Kolesnikov et al. // Eurasian Soil Science. — 2025. — № 58 (10). — P. 136. — DOI: 10.1134/S1064229325601106
2. Ob utverzhdenii normativov kachestva vody' vodny'x ob'ektov ry'boxozajstvennogo znacheniya, v tom chisle normativov predel'no dopustimy'x koncentracij zagryaznyayushhix veshhestv v vodax vodny'x ob'ektov ry'boxozajstvennogo znacheniya : Prikaz Federal'nogo agentstva po ry'bologstvu ot 26 maya 2025 g. № 296 [On approval of water quality standards for fishery water bodies, including maximum permissible concentrations of pollutants in the waters of fishery water bodies: Order of the Federal Agency for Fisheries No. 296 of May 26, 2025] // Official Internet Portal of Legal Information. — 2025. — URL: <http://publication.pravo.gov.ru/document/0001202506020069>. (accessed: 24.08.26) [in Russian]
3. Sazonov A.A. Sinkhronnye issledovaniya kachestva vody reki Volgi [Synchronous studies of water quality in the Volga River] / A.A. Sazonov, A.A. Lisina, O.N. Erina [et al.] // Vodnoe khozyaystvo Rossii: problemy, tekhnologii, upravlenie [Water Sector of Russia: Problems, Technologies, Management]. — 2024. — № 5. — P. 98–114. — DOI: 10.35567/19994508-2024-5-98-114 [in Russian]
4. Udalov I.V. Gidrokhimicheskaya kharakteristika poverkhnostnykh i gruntovykh vod Lisichanskogo i Almazno-Mar'evskogo geologo-promyshlennykh rayonov Severo-Vostochnogo Donbassa [Hydrochemical characteristics of surface and ground waters of the Lysichansk and Almazno-Marievka geological and industrial regions of the North-Eastern Donbass] / I.V. Udalov // Journal of Geology, Geography and Geoecology. — 2014. — Vol. 22. — № 3 (2). — P. 2–11. [in Russian]
5. Khoroshevskaya V.O. Sravnitel'naya kharakteristika mikroelementnogo sostava rechnykh vod, dreniruyushchikh Donetskii i Kuzbasskii ugolnye basseyny [Comparative characteristics of trace element composition of river waters draining the Donetsk and Kuznetsk coal basins] / V.O. Khoroshevskaya // Mezhdunarodny nauchno-issledovatel'sky zhurnal [International Research Journal]. — 2022. — № 5-1 (119). — P. 124–134. — DOI: 10.23670/IRJ.2022.119.5.062 [in Russian]
6. Reutova N.V. Prirodnoe i antropogennoe zagryaznenie mikroelementami verkhoviy basseyna r. Ardon [Natural and anthropogenic pollution with trace elements in the upper reaches of the Ardon River basin] / N.V. Reutova, T.V. Reutova, F.R. Dreeva // Geologiya i geofizika Yuga Rossii [Geology and Geophysics of Russian South]. — 2025. — Vol. 15. — № 2. — P. 219–230. — DOI: 10.46698/VNC.2025.46.62.001 [in Russian]



7. Batley G. E. Metal contaminants of emerging concern in aquatic systems / G. E. Batley, P. G. C. Campbell // Environmental Chemistry. — 2022. — Vol. 19. — № 1. — P. 23–40. — DOI: 10.1071/EN22030
8. Sawasdee B. Metal sensitivity of the embryonic development of the ramshorn snail *Marisa cornuarietis* (Prosobranchia) / B. Sawasdee, H. R. Köhler // Ecotoxicology. — 2010. — Vol. 19. — № 8. — P. 1487–1495. — DOI: 10.1007/s10646-010-0534-8
9. Osborne R. K. Development of an embryo toxicity test to assess the comparative toxicity of metal exposure on different life stages of freshwater gastropods / R. K. Osborne, M. M. Ijzerman, C. Venier [et al.] // Environmental Toxicology and Chemistry. — 2023. — Vol. 42. — № 8. — P. 1791–1805. — DOI: 10.1002/etc.5686
10. Calabrese E. J. Lithium and hormesis: Enhancement of adaptive responses and biological performance via hormetic mechanisms / E. J. Calabrese, P. Pressman, A. W. Hayes [et al.] // Journal of Trace Elements in Medicine and Biology. — 2023. — Vol. 78. — P. 127156. — DOI: 10.1016/j.jtemb.2023.127156
11. Noguera L. Integration of responses to lithium in mussels at different levels of biological complexity / L. Noguera, D. Benito, T. Briaudeau [et al.] // Chemosphere. — 2025. — Vol. 371. — P. 144021. — DOI: 10.1016/j.chemosphere.2024.144021.
12. Campos S. Behavioural and biochemical responses of the sea snail *Tritia reticulata* to lithium concentration gradient / S. Campos, C. Leite, J. Pinto [et al.] // Aquatic Toxicology. — 2023. — Vol. 261. — P. 106629. — DOI: 10.1016/j.aquatox.2023.106629.